arXiv:1503.03418v1 [math.NT] 11 Mar 2015

Note on super congruences modulo p^2

Zhi-Hong Sun

School of Mathematical Sciences
Huaiyin Normal University
Huaian, Jiangsu 223001, P.R. China
Email: zhihongsun@yahoo.com
Homepage: http://www.hytc.edu.cn/xsjl/szh

Abstract

Let p be an odd prime, and let m be an integer with $p\nmid m$. In this paper show that

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}}{m^k}\equiv 0\pmod p\quad\text{implies}\quad\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}}{m^k}\equiv 0\pmod {p^2}.$$

MSC: Primary 11A07, Secondary 33C45, 05A19
Keywords: congruence; Legendre polynomial

1. Introduction

Let $[x]$ be the greatest integer not exceeding x. For a prime p let $\mathbb{Z}_p$ be the set of rational numbers whose denominator is not divisible by p. Let $\{P_n(x)\}$ be the Legendre polynomials given by

$$P_0(x)=1,\ P_1(x)=x,\ (n+1)P_{n+1}(x)=(2n+1)xP_n(x)-nP_{n-1}(x)\ (n\ge 1).$$

It is well known that (see [MOS, pp. 228-232], [B1] and [B2])

$$(1.1)\qquad P_n(x)=\frac{1}{2^n}\sum_{k=0}^{[n/2]}\binom nk(-1)^k\binom{2n-2k}{n}x^{n-2k}=\sum_{k=0}^{n}\binom nk\binom{n+k}{k}\Big(\frac{x-1}{2}\Big)^k.$$

From (1.1) we see that $P_n(-x)=(-1)^nP_n(x)$.

Let $p>3$ be a prime. In 2003 Rodriguez-Villegas [RV] conjectured that

$$(1.2)\qquad \begin{aligned}&\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{3k}{k}}{108^k}\equiv 0\ (\text{mod }p^2)\quad\text{for}\quad p\equiv 2\ (\text{mod }3),\\ &\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{4k}{2k}}{256^k}\equiv 0\ (\text{mod }p^2)\quad\text{for}\quad p\equiv 5,7\ (\text{mod }8),\\ &\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{3k}{k}\binom{6k}{3k}}{1728^k}\equiv 0\ (\text{mod }p^2)\quad\text{for}\quad p\equiv 3\ (\text{mod }4).\end{aligned}$$

In 2005 Mortenson [M] proved these congruences modulo p, in 2012 Z.W. Sun [Su3] confirmed (1.2). Motivated by Mortenson's work, in [Su1,Su2] Z.W. Sun posed many conjectures concerning the following sums modulo p^2:

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m^k},\ \sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{3k}{k}}{m^k},\ \sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{4k}{2k}}{m^k}\quad\text{and}\quad\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{3k}{k}\binom{6k}{3k}}{m^k},$$

where m is an integer with $p\nmid m$. In [S2-S5] the author solved some of his conjectures by establishing the following congruences:

$$(1.3)\quad \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m^k}\equiv P_{\frac{p-1}{2}}\Big(\sqrt{1-\frac{64}{m}}\Big)^2 \pmod{p^2},$$

$$(1.4)\quad \sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{3k}{k}(x(1-27x))^k\equiv\Big(\sum_{k=0}^{p-1}\binom{2k}{k}\binom{3k}{k}x^k\Big)^2 \pmod{p^2},$$

$$(1.5)\quad \sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{4k}{2k}(x(1-64x))^k\equiv\Big(\sum_{k=0}^{p-1}\binom{2k}{k}\binom{4k}{2k}x^k\Big)^2 \pmod{p^2},$$

$$(1.6)\quad \sum_{k=0}^{p-1}\binom{2k}{k}\binom{3k}{k}\binom{6k}{3k}(x(1-432x))^k\equiv\Big(\sum_{k=0}^{p-1}\binom{2k}{k}\binom{3k}{k}\binom{6k}{3k}x^k\Big)^2 \pmod{p^2}.$$

It is easily seen that (see [S1, pp.1916-1917, 1920], [S3, p.1953] and [S5, p.182])

$$(1.7)\quad \begin{aligned}&\binom{-\frac12}{k}^2=\frac{\binom{2k}{k}^2}{16^k},\quad \binom{-\frac13}{k}\binom{-\frac23}{k}=\frac{\binom{2k}{k}\binom{3k}{k}}{27^k},\\ &\binom{-\frac14}{k}\binom{-\frac34}{k}=\frac{\binom{2k}{k}\binom{4k}{2k}}{64^k},\quad \binom{-\frac16}{k}\binom{-\frac56}{k}=\frac{\binom{3k}{k}\binom{6k}{3k}}{432^k}.\end{aligned}$$

Let p be an odd prime, $a\in\mathbb{Z}_p$ and let $\langle a\rangle_p\in\{0,1,\dots,p-1\}$ be given by $a\equiv\langle a\rangle_p \pmod p$. In [S6], the author proved that

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}}{4^k}\equiv 0 \pmod{p^2}\quad\text{for}\quad \langle a\rangle_p\equiv 1 \pmod 2.$$

Let $(a)_k=a(a+1)\cdots(a+k-1)=(-1)^k k!\binom{-a}{k}$ and

$${}_rF_s\left(\begin{matrix}a_1,\dots,a_r\\ b_1,\dots,b_s\end{matrix}\,\middle|\,z\right)=1+\sum_{k=1}^{\infty}\frac{(a_1)_k\cdots(a_r)_k}{(b_1)_k\cdots(b_s)_k}\cdot\frac{z^k}{k!}.$$

After reading the author's preprint (arXiv:1101.1050) involving (1.4)-(1.6), in the email to the author on January 11, 2011 Wadim Zudilin wrote: "It's probably worth mentioning that the proofs of Theorems 2.1, 2.4 and possibly Theorem 3.1 from your arXiv:1101.1050 can be simplified. Your congruences assume the form

$$\left(\sum_{k=0}^{p-1}\frac{(a)_k(1-a)_k}{k!^2}x^k\right)^2\equiv\sum_{k=0}^{p-1}\frac{(a)_k(1-a)_k}{k!^2}\binom{2k}{k}(x(1-x))^k \pmod{p^2},$$

where $a = 1/3,\ 1/4$ or $1/6$ ($1/2$ is possible as well). Note that this follows from the identity

$$\left(\sum_{k=0}^{\infty}\frac{(a)_k(1-a)_k}{k!^2}x^k\right)^2=\sum_{k=0}^{\infty}\frac{(a)_k(1-a)_k}{k!^2}\binom{2k}{k}(x(1-x))^k$$

truncated to p terms. If $k > p/2$, then the coefficients on both sides are $0 \pmod{p^2}$ (as you already used in the proof, e.g., of Theorem 2.1)." In the email on January 15, 2011 Zudilin wrote: The identity is a combination of the Gauss quadratic transformation ([Ba, p.88, Eq. (2)])

$${}_2F_1\left(\begin{matrix}A,\ B\\ A+B+\frac{1}{2}\end{matrix}\,\middle|\, 4z(1-z)\right)={}_2F_1\left(\begin{matrix}2A,\ 2B\\ A+B+\frac{1}{2}\end{matrix}\,\middle|\, z\right),$$

and Clausen's original identity ([Sl, p.75, Eq.(2.5.7)])

$${}_2F_1\left(\begin{matrix}a,\ b\\ a+b+\frac{1}{2}\end{matrix}\,\middle|\, z\right)^2={}_3F_2\left(\begin{matrix}2a,\ 2b,\ a+b\\ a+b+\frac{1}{2},\ 2a+2b\end{matrix}\,\middle|\, z\right).$$

Let p be an odd prime and $a \in \mathbb{Z}_p$. In this note, inspired by Zudilin's comments we give an elementary proof of the following generalization of (1.4)-(1.6):

$$\Big(\sum_{k=0}^{p-1}\binom{a}{k}\binom{-1-a}{k}x^k\Big)^2\equiv\sum_{k=0}^{p-1}\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}(x(1-x))^k \pmod{p^2}.$$

As an application, for $m \in \mathbb{Z}_p$ with $m \not\equiv 0 \pmod p$ we show that

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}}{m^k}\equiv 0 \pmod p \quad\text{implies}\quad \sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}}{m^k}\equiv 0 \pmod{p^2}.$$

2. Main results

Lemma 2.1 ([S3, Lemma 3.2]). *For any nonnegative integer n we have*

$$P_n(\sqrt{1+4x})^2=\sum_{k=0}^{n}\binom{n}{k}\binom{n+k}{k}\binom{2k}{k}x^k.$$

As Zudilin noted, Lemma 2.1 can also be deduced from Clausen's identity, Gauss' quadratic transformation for hypergeometic series and (1.1).

Theorem 2.1. *Let p be an odd prime and $a \in \mathbb{Z}_p$. Then*

$$\sum_{k=0}^{p-1}\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}x^k\equiv P_{\langle a\rangle_p}(\sqrt{1-4x})^2\equiv P_{p-1-\langle a\rangle_p}(\sqrt{1-4x})^2 \pmod p.$$

Proof. By Lemma 2.1,

$$P_{\langle a\rangle_p}(\sqrt{1-4x})^2=\sum_{k=0}^{\langle a\rangle_p}\binom{2k}{k}\binom{\langle a\rangle_p}{k}\binom{\langle a\rangle_p+k}{k}(-x)^k$$

$$\begin{aligned}&=\sum_{k=0}^{p-1}\binom{2k}{k}\binom{\langle a\rangle_p}{k}\binom{\langle a\rangle_p+k}{k}(-x)^k\\&\equiv\sum_{k=0}^{p-1}\binom{2k}{k}\binom{a}{k}\binom{a+k}{k}(-x)^k\\&=\sum_{k=0}^{p-1}\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}x^k \pmod p.\end{aligned}$$

To complete the proof, we note that $P_{p-1-\langle a\rangle_p}(t)\equiv P_{\langle a\rangle_p}(t) \pmod p$ by [S4, Lemma 2.2].

Remark 2.1 In the cases $a=-\frac12,-\frac13,-\frac14,-\frac16$, Theorem 2.2 was given by the author in [S2, Theorem 4.1], [S3, Theorems 3.2 and 4.2] and [S5, Theorem 4.1].

Corollary 2.1. *Let p be an odd prime and $a,x\in\mathbb{Z}_p$. Then $\sum_{k=0}^{p-1}\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}x^k\equiv 0 \pmod p$ implies*

$$P_{\langle a\rangle_p}(\sqrt{1-4x})\equiv P_{p-1-\langle a\rangle_p}(\sqrt{1-4x})\equiv 0 \pmod p.$$

Proof. By (1.1),

$$P_n(\sqrt{1-4x})=\frac{1}{2^n}(\sqrt{1-4x})^{n-2[\frac n2]}\sum_{k=0}^{[n/2]}\binom{n}{k}(-1)^k\binom{2n-2k}{n}(1-4x)^{[\frac n2]-k}.$$

Thus $P_n(\sqrt{1-4x})^2\equiv 0 \pmod p$ implies $P_n(\sqrt{1-4x})\equiv 0 \pmod p$. Now applying Theorem 2.1 we deduce the result.

Lemma 2.2. *For any nonnegative integer n we have*

$$\sum_{k=0}^{n}\binom{a}{k}\binom{-1-a}{k}\binom{a}{n-k}\binom{-1-a}{n-k}=\sum_{k=0}^{n}\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}\binom{k}{n-k}(-1)^{n-k}.$$

Proof. Let $S_1(n)$ and $S_2(n)$ be the sums on the left and right hands of the identity, respectively. Using Maple and the Zeilberger algorithm we find that for $i=1,2$,

$$n^3S_i(n)=(2n-1)(n^2-n-2a(a+1))S_i(n-1)+(n-1)(2a+n)(2a+2-n)S_i(n-2)\ (n\ge 2).$$

Since $S_1(0)=1=S_2(0)$ and $S_1(1)=-2a(a+1)=S_2(1)$, applying the above we deduce that $S_1(n)=S_2(n)$.

Remark 2.2 In the cases $a=-\frac13,-\frac14,-\frac16$, the identity was given by the author in [S2,S3,S5].

Theorem 2.2. *Let p be an odd prime and $a\in\mathbb{Z}_p$. Then*

$$\Big(\sum_{k=0}^{p-1}\binom{a}{k}\binom{-1-a}{k}x^k\Big)^2\equiv\sum_{k=0}^{p-1}\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}(x(1-x))^k \pmod{p^2}.$$

Taking $a=-\frac13,-\frac14,-\frac16$ in Theorem 2.2 and then applying (1.7) we get (1.4)-(1.6). Thus, Theorem 2.2 is a generalization of (1.4)-(1.6).

Proof of Theorem 2.2. For $k\in\{\frac{p+1}{2},\dots,p-1\}$ we see that $p\mid\binom{2k}{k}$ and

$$\binom{a}{k}\binom{-1-a}{k}=(-1)^k\binom{a}{k}\binom{a+k}{k}$$

$$= (-1)^k \frac{(a+k)(a+k-1)\cdots(a-k+1)}{k!} \equiv 0 \pmod p.$$

Thus, applying Lemma 2.2 we see that

$$\begin{aligned}
&\sum_{k=0}^{p-1} \binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}(x(1-x))^k \\
&\equiv \sum_{k=0}^{(p-1)/2} \binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}(x(1-x))^k \\
&= \sum_{k=0}^{(p-1)/2} \binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}x^k \sum_{r=0}^{k}\binom{k}{r}(-x)^r \\
&= \sum_{n=0}^{p-1} x^n \sum_{k=0}^{\min\{n,\frac{p-1}{2}\}} \binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}\binom{k}{n-k}(-1)^{n-k} \\
&\equiv \sum_{n=0}^{p-1} x^n \sum_{k=0}^{n} \binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}\binom{k}{n-k}(-1)^{n-k} \\
&= \sum_{n=0}^{p-1} x^n \sum_{k=0}^{n} \binom{a}{k}\binom{-1-a}{k}\binom{a}{n-k}\binom{-1-a}{n-k} \\
&= \sum_{k=0}^{p-1} \binom{a}{k}\binom{-1-a}{k}x^k \sum_{n=k}^{p-1}\binom{a}{n-k}\binom{-1-a}{n-k}x^{n-k} \\
&= \Big(\sum_{k=0}^{p-1}\binom{a}{k}\binom{-1-a}{k}x^k\Big)\Big(\sum_{r=0}^{p-1}\binom{a}{r}\binom{-1-a}{r}x^r - \sum_{r=p-k}^{p-1}\binom{a}{r}\binom{-1-a}{r}x^r\Big) \pmod{p^2}.
\end{aligned}$$

For $0 \le k \le p-1$ we see that $p \mid \binom{a}{k}$ for $k > \langle a\rangle_p$. For $0 \le k \le \langle a\rangle_p$ and $p-k \le r \le p-1$ we see that $r \ge p-k > p-1-\langle a\rangle_p$ and so $p \mid \binom{-1-a}{r}$. Hence for $0 \le k \le p-1$ and $p-k \le r \le p-1$ we have $p \mid \binom{a}{k}\binom{-1-a}{r}$, $p \mid \binom{-1-a}{k}\binom{a}{r}$ and so $p^2 \mid \binom{a}{k}\binom{-1-a}{k}\binom{a}{r}\binom{-1-a}{r}$. Now combining all the above we deduce the result.

Theorem 2.3. *Let p be an odd prime and $m \in \mathbb{Z}_p$ with $m \not\equiv 0 \pmod p$. Then*

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}}{m^k} \equiv 0 \pmod p \quad \textit{implies} \quad \sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}}{m^k} \equiv 0 \pmod{p^2}.$$

Proof. Suppose $\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}}{m^k} \equiv 0 \pmod p$. By Corollary 2.1, $P_{\langle a\rangle_p}(\sqrt{1-\frac{4}{m}}) \equiv 0 \pmod p$. Thus,

$$\begin{aligned}
&\sum_{k=0}^{p-1}\binom{a}{k}\binom{-1-a}{k}\Big(\frac{1-\sqrt{4/m}}{2}\Big)^k \\
&\equiv \sum_{k=0}^{p-1}\binom{\langle a\rangle_p}{k}\binom{-1-\langle a\rangle_p}{k}\Big(\frac{1-\sqrt{4/m}}{2}\Big)^k
\end{aligned}$$

$$= \sum_{k=0}^{\langle a\rangle_p}\binom{\langle a\rangle_p}{k}\binom{\langle a\rangle_p+k}{k}\Big(\frac{\sqrt{4/m}-1}{2}\Big)^k$$

$$= P_{\langle a\rangle_p}\Big(\sqrt{1-\frac{4}{m}}\Big)\equiv 0 \pmod p.$$

By Theorem 2.2,

$$\Big(\sum_{k=0}^{p-1}\binom{a}{k}\binom{-1-a}{k}\Big(\frac{1-\sqrt{4/m}}{2}\Big)^k\Big)^2\equiv\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{a}{k}\binom{-1-a}{k}}{m^k} \pmod {p^2}.$$

Thus the result follows.

Taking $a=-\frac12,-\frac13,-\frac14,-\frac16$ in Theorem 2.3 and then applying (1.7) we deduce the following result.

Corollary 2.2. *Let p be an odd prime, and $m\in\mathbb{Z}_p$ with $m\not\equiv 0 \pmod p$. Then*

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m^k}\equiv 0 \pmod p \quad \textit{implies} \quad \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m^k}\equiv 0 \pmod {p^2},$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{3k}{k}}{m^k}\equiv 0 \pmod p \quad \textit{implies} \quad \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{3k}{k}}{m^k}\equiv 0 \pmod {p^2},$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{4k}{2k}}{m^k}\equiv 0 \pmod p \quad \textit{implies} \quad \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{4k}{2k}}{m^k}\equiv 0 \pmod {p^2}$$

and

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{3k}{k}\binom{6k}{3k}}{m^k}\equiv 0 \pmod p \quad \textit{implies} \quad \sum_{k=0}^{p-1}\frac{\binom{2k}{k}\binom{3k}{k}\binom{6k}{3k}}{m^k}\equiv 0 \pmod {p^2}.$$

We remark that Corollary 2.2 can be easily deduced from [S2, Theorem 4.1], [S3, Theorems 3.2 and 4.2] and [S5, Theorem 4.1].

Theorem 2.4. *Let p be an odd prime and $u\in\mathbb{Z}_p$.*

(i) *If $u\not\equiv\frac14,\frac1{16} \pmod p$ and $\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{3k}{k}(\frac{u^2}{(1-4u)^3})^k\equiv 0 \pmod p$, then*

$$\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{3k}{k}\Big(\frac{-u}{(1-16u)^3}\Big)^k\equiv 0 \pmod {p^2}.$$

(ii) *If $u\not\equiv-\frac13,-\frac1{27} \pmod p$ and $\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{4k}{2k}(\frac{u^3}{(1+3u)^4})^k\equiv 0 \pmod p$, then*

$$\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{4k}{2k}\Big(\frac{u}{(1+27u)^4}\Big)^k\equiv 0 \pmod {p^2}.$$

Proof. This is immediate from Corollary 2.2 and [S7, Corollaries 3.1 and 5.2].

Corollary 2.3. *Let $p>3$ be a prime. Then*

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{3k}{k}}{1458^k}\equiv 0 \pmod {p^2} \quad \textit{for} \quad p\equiv 5 \pmod 6$$

and

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{3k}{k}}{15^{3k}}\equiv 0 \pmod{p^2}\quad \textit{for}\quad p\equiv 11,14 \pmod{15}.$$

Proof. Taking $u=-\frac{1}{2}$ in Theorem 2.4(i) and then applying (1.2) we obtain the first congruence. Taking $u=1$ in Theorem 2.4(i) and then applying [S2, Theorems 4.6] we obtain the second congruence.

Remark 2.3 Let p be a prime of the form $6k+5$. In [S7] the author proved that $\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{3k}{k}}{1458^k}\equiv 0 \pmod p$ and conjectured that $\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2\binom{3k}{k}}{1458^k}\equiv 0 \pmod{p^3}$.

Acknowledgment

The author is indebted to Wadim Zudilin for his contribution to Theorem 2.2. See comments in Section 1.

References

[Ba] W. N. Bailey, *Generalized hypergeometric series*, Cambridge Math. Tracts **32** Cambridge Univ. Press, Cambridge ,1935; 2nd reprinted edition, Stechert-Hafner, New York–London, 1964.

[B1] H. Bateman, *Higher Transcendental Functions, Vol.I*, McGraw-Hill, New York, 1953.

[B2] H. Bateman, *Higher Transcendental Functions, Vol.II*, McGraw-Hill, New York, 1953.

[MOS] W. Magnus, F. Oberhettinger and R.P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics* (*3rd ed.*), Springer-Verlag, New York, 1966, pp. 25-32.

[M] E. Mortenson, *Supercongruences for truncated* ${}_{n+1}F_n$ *hypergeometric series with applications to certain weight three newforms*, Proc. Amer. Math. Soc. **133**(2005), 321-330.

[RV] F. Rodriguez-Villegas, *Hypergeometric families of Calabi-Yau manifolds* in: Noriko Yui, James D. Lewis (Eds.), Calabi-Yau Varieties and Mirror Symmetry , Toronto, ON, 2001, in: Fields Inst. Commun., vol. 38, Amer. Math. Soc., Providence, RI, 2003, pp.223-231.

[Sl] L. J. Slater, *Generalized hypergeometric functions*, Cambridge Univ. Press, Cambridge, 1966.

[S1] Z. H. Sun, *Congruences concerning Legendre polynomials*, Proc. Amer. Math. Soc. **139**(2011), 1915-1929.

[S2] Z. H. Sun, *Congruences involving* $\binom{2k}{k}^2\binom{3k}{k}$, J. Number Theory **133**(2013), 1572-1595.

[S3] Z. H. Sun, *Congruences concerning Legendre polynomials II*, J. Number Theory **133**(2013), 1950-1976.

[S4] Z. H. Sun, *Congruences concerning Legendre polynomials III*, Int. J. Number Theory **9**(2013), 965-999.

[S5] Z. H. Sun, *Legendre polynomials and supercongruences*, Acta Arith. **159**(2013), 169-200.

[S6] Z. H. Sun, *Generalized Legendre polynomials and related supercongruences*, J. Number Theory **143**(2014), 293-319.

[S7] Z. H. Sun, *Congruences for Domb and Almkvist-Zudilin numbers*, arXiv:1302.1715v7, 2015.

[Su1] Z.W. Sun, *Open conjectures on congruences*, arXiv:0911.5665v59, 2011.

[Su2] Z.W. Sun, *Super congruences and Euler numbers*, Sci. China Math. **54**(2011), 2509-2535.

[Su3] Z. W. Sun, *On sums involving products of three binomial coefficients*, Acta Arith. **156**(2012), 123-141.